

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/110865/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Jones, Benjamin, Cullen-Unsworth, Leanne C. ORCID: <https://orcid.org/0000-0002-9134-7266>, Howard, Robert and Unsworth, Richard K.F. 2018. Complex yet fauna-deficient seagrass ecosystems at risk in southern Myanmar. *Botanica Marina* 61 (3) , pp. 193-204. 10.1515/bot-2017-0082 file

Publishers page: <http://dx.doi.org/10.1515/bot-2017-0082>
<<http://dx.doi.org/10.1515/bot-2017-0082>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 Complex yet fauna-deficient seagrass ecosystems at risk in southern Myanmar

2 Benjamin L. Jones^{1,2*}, Leanne C. Cullen-Unsworth^{1,2}, Robert Howard³ & Richard K.F.
3 Unsworth^{2,4}

4
5 ¹Sustainable Places Research Institute, Cardiff University, 33 Park Place, Cardiff, CF10 3BA,
6 UK

7 ²Project Seagrass, 33 Park Place, Cardiff, UK

8 ³Fauna & Flora International, Yangon, Myanmar

9 ⁴Seagrass Ecosystem Research Group, College of Science, Swansea University, Swansea, SA2
10 8PP, UK

11
12 *Corresponding Author: JonesB67@cardiff.ac.uk

14 **Abstract**

15 Dependence on seafood across Southeast Asia is extensive. Myanmar is no exception,
16 but the country's provisioning marine ecosystems are threatened. Seagrass is one habitat that
17 is frequently overlooked in management as an important fisheries resource, despite its nursery
18 function. In Myanmar, research on seagrass habitats is particularly sparse, and as a result, our
19 understanding of seagrass exploitation remains limited. In this study, we provide a baseline
20 assessment of the seagrass-associated fish assemblages at four locations in the Myeik
21 Archipelago in southern Myanmar using mono Baited Remote Underwater Video systems.
22 Across the sites surveyed only 12 taxa of motile fauna were recorded. Relative to other regional
23 and global studies this figure is meagre. Our data adds to a growing literature suggesting that
24 the marine habitats of Myanmar are in decline. Despite the lack of recorded seagrass associated
25 fauna, our study revealed minimal impacts to seagrass meadows from eutrophication or
26 sedimentation, and the meadows included appeared to be healthy. The sites with the highest
27 number of motile fauna were within Myanmar's only National Marine Park offering some
28 optimism for the effectiveness of protection, but further assessments are required to allow
29 targeted management of Myanmar's seagrass meadows.

30
31 **Keywords:** Seagrass fisheries, Myanmar, Myeik Archipelago, over-fishing, fisheries
32 resources, BRUV

Introduction

Dependence on seafood across Southeast Asia is extensive (Donner and Potere, 2007), and this holds true for Myanmar (Russell, 2015). However, tropical marine ecosystems are in decline (Burke et al., 2011; Giri et al., 2011; Pauly, 1998; Waycott et al., 2009), along with their associated fisheries (Pauly, 1998). While there is a magnitude of reasons for fishery decline, Malthusian over-fishing has been pivotal (McClanahan et al., 2009). In Myanmar, due to historical political uncertainty, formal management of marine ecosystems has been absent until relatively recently.

Tropical marine ecosystems, like those present along the 3000km coastline of Myanmar, form an inter-connected seascape that supports a diversity of marine organisms throughout their life stages. Connected coastal habitats provide daily foraging for adult fish to residency areas for juveniles (Harborne et al., 2006; Harborne et al., 2008; Nagelkerken et al., 2008; Unsworth et al., 2008). Seagrass meadows are one component of these tropical marine systems, also providing critical fishery grounds (de la Torre-Castro and Ronnback, 2004; Exton, 2010; Nordlund et al., 2017; Unsworth and Cullen, 2010). Seagrass meadows provide nursery habitat for numerous fish species that are relied upon for food and livelihoods (Beck et al., 2001; Gillanders, 2006; Heck et al., 2003; Nagelkerken et al., 2012; Unsworth et al., 2014a).

Across the Indo-Pacific, marine research, monitoring and conservation funding has been geared towards coral reefs and to a lesser extent mangroves. This trend is mostly a result of tropical coastal marine fisheries often being referred to misleadingly as ‘coral reef fisheries’ (Nordlund et al., 2017; Unsworth and Cullen, 2010). Despite recognition for their valuable ecosystem services, the role of seagrasses in supporting fisheries productivity and importantly food supply has largely been ignored until recently (de la Torre-Castro et al., 2014; Duarte et al., 2008; Nordlund et al., 2017; Unsworth and Cullen, 2010). In Myanmar, limited research effort means we have a more limited understanding of seagrass ecosystems. To date, seagrass ecosystem research in Myanmar has focused on mapping meadows and recording the seagrass species’ present (Giardino et al., 2016; Ilangakoon and Tun, 2007; Novak et al., 2009; Soe-Htun et al., 2015; Soe-Htun et al., 2001). In 2001 Soe-Htun et al. (2001), described seagrasses in Myanmar as being in “pristine and climax conditions” with “no stresses.” In 2015, however, the same authors suggested that this is no longer the case (Soe-Htun et al., 2015). Seagrass meadows in Myanmar are now suffering from the same regional problems, including eutrophication and sedimentation, observed elsewhere in Southeast Asia (Ooi et al., 2011; Satumanatpan, 2008; Satumanatpan et al., 2011). Previous monitoring reports suggest seagrass

meadows in Myanmar support a rich biodiversity of marine life, including fish from families including Chaetodontidae (butterflyfishes), Pomacanthidae (angelfishes), Labridae (wrasses), Siganidae (rabbitfishes), Mugilidae (mulletts) and Clupeidae (herring) (Soe-Htun et al., 2001). Surveys of Myanmar's coral reef and demersal fisheries in recent years have shown a rapid decline in biodiversity (Howard et al., 2014; Obura et al., 2014; Russell, 2015, 2016). Therefore, it is likely that seagrass fisheries have also seen a similar fate.

Understanding habitat links to fisheries, while critical for short-term fisheries management, is important for understanding the vulnerability, and resilience of marine systems to change (Folke, 2006; McClanahan et al., 2009). There is an urgent need to understand the role that different habitat types have in supporting tropical marine fisheries within the Indo-Pacific region. Given the growing evidence of the role, that seagrass meadows play in supporting Indo-Pacific marine fisheries (Nordlund et al., 2017), here we provide a baseline assessment of the seagrass meadows and their associated fish assemblages at four locations in the Myeik Archipelago in southern Myanmar and discuss the implications of our findings.

Methods

Study location

The Myeik Archipelago (formerly Mergui) is comprised of around 800 islands covering around 36,000 km² in the southern Tanintharyi coastal region of Myanmar (Fig. 1). The Archipelago is inhabited by a population of approximately 2000-3000 semi-nomadic people, the Moken (also referred to as Sea Gypsies or *Salone* in Burmese) (Schneider et al., 2014). During April and May 2016 floral and faunal assessments were conducted within seagrass meadows at four sites as follows: Taw Wet North (11.41°, 98.12°); Lampi East (10.70°, 98.28°); Bo Cho (10.67°, 98.26°) and; Nyaung Wee (10.50°, 98.23°). Taw Wet North was the northernmost site used in this study, and the seagrass meadow was located in front of a small mangrove habitat, with a small freshwater input. Seagrass meadows at Lampi East and Bo Cho were both located in front of a sandy beach in the absence of mangrove. Lampi East and Bo Cho were within the Lampi Marine National Park (MNP) (Table 1). Lampi Island MNP includes around 3000 people living in 5 settlements within the boundary of the park (MOECAP and Oikos, 2015). Although not within the park, Nyaung Wee is included in this population estimate and was the southernmost site. The seagrass meadow at Nyaung Wee is situated in front of a small mangrove habitat. At all sites, the substrate was muddy sand nearshore, becoming sandy mud further offshore (McKenzie et al., 2001).

Seagrass morphometrics

At each sampling site, 34 haphazardly placed 0.25 m² quadrats were sampled from within the seagrass meadow at low tide. Shoot density (0.0225 m²) was recorded as was total percentage cover and floral species composition (McKenzie et al., 2001). Canopy height was also recorded using the mean height of three leaves in each quadrat. Percentage epiphyte and algal cover were recorded using the Seagrass-Watch quadrat metrics (McKenzie et al., 2001).

Biodiversity assessments

The relative abundance and diversity of fish assemblages were assessed using mono-camera Baited Remote Underwater Video systems (BRUVs). Fish were identified to species level where possible. The mono-BRUVs were constructed based on designs by Cappo et al. (2004), using a stainless steel tripod-style frame as a mount for a GoPro Hero 4 camera. A bait arm (20 mm stainless steel conduit) extending 1m from the base plate of the camera supported a plastic bait container (112 cm³), holding standardised bait (ground goatfish and sardine – sourced locally), which was replenished before every deployment.

Five sets of three deployments, spaced 50m apart (15 samples) were conducted at Taw Wet North and four sets of three deployments, again spaced 50m apart (12 samples) were conducted at Lampi East, Bo Cho and Nyaung Wee. BRUVs were deployed for 30 minutes which is considered adequate time to assess fish assemblages while remaining cost-effective (Haggitt et al., 2014; Kelaher et al., 2014; Malcolm et al., 2015; Wraith et al., 2013; Wraith, 2007). Additionally, a short sampling duration enables a higher number of samples to be collected, achieving a great spatial representation of the variability of the fish assemblages (Unsworth et al., 2014b). BRUVs were deployed at depths of 0.5 to 1.5m on an incoming tide. All BRUVs sampling was conducted during daylight hours.

Video footage was analysed to determine the MaxN of each fish species in each sample. MaxN is a metric commonly used for the quantification of the relative abundance of fish observed on underwater video (Cappo et al., 2004; Unsworth et al., 2014b). MaxN counts the maximum number of fish recorded at any one time (single video frame) and therefore removes concerns associated with double counting of individual fish (Priede et al., 1994). All footage was analysed using the specialised SeaGIS software EventMeasure v.3.51. MaxN was determined for each species in every video frame throughout the 30 minutes of footage. The highest MaxN for each species at the end of each 30 minutes was then used in further analysis.

Data analysis

Data were tested for homogeneity of variance and normality. Where data were not normal, log transformations were performed so that data met the assumptions of parametric tests. One-way ANOVA was used to test for differences in the key seagrass morphometrics across sites with Bonferroni post-hoc tests for differences between sites using the software SPSS v.23. Analysis of differences in the structure of fish assemblages between sites was conducted using multivariate non-metric multidimensional scaling ordination (nMDS) in PRIMER v.6.1.5, and a 2-way analysis of similarities (ANOSIM) was used to investigate differences identified from MDS (Clarke and Warwick, 1994). All summary data are presented as means \pm standard deviation.

Results

Seagrass condition

All seagrass meadows surveyed were mixed species meadows. The dominant species across all sites was *Cymodocea rotundata* (>80%; Table 2). Seagrass morphometrics differed across the four sites (see Fig. 2). Significant differences in seagrass cover were observed between sites ($F_{3, 125} = 4.4250$, $p < 0.001$). Highest percentage cover values were recorded at Nyaung Wee (48.9 ± 27.5 %), which were significantly different from the lowest values at Bo Cho (28.3 ± 21.1 %). Seagrass cover at Taw Wet North (32.2 ± 25.5 %) and Lampi East (42.6 ± 28.7 %) did not significantly differ from the other two sites. There were no significant differences in shoot density across sites, where highest values were recorded at Lampi East ($266.4 \pm 155.13 \text{ m}^{-2-1}$) and lowest at Taw Wet North ($219.2 \pm 146.1 \text{ m}^{-2-1}$).

Significant differences in canopy height were recorded between sites ($F_{3, 125} = 7.231$, $p < 0.001$). Values were highest at Taw Wet North (9.9 ± 2.8 cm) and lowest at Bo Cho (6.6 ± 2.8 cm). Significant differences in canopy height were observed between Taw Wet North and Bo Cho, and Taw Wet North and Lampi East (7.5 ± 6.6 cm). Significant differences in epiphyte cover ($F_{3, 125} = 11.635$, $p < .001$) were observed between sites. Epiphyte cover was characteristically low at Taw Wet North, Bo Cho and Nyaung Wee and high values at Lampi East (20.3 ± 23.3 %) were responsible for differences between sites. Taw Wet North had no algae present and was responsible for significant differences in algae cover between sites ($F_{3, 125} = 7.602$, $p < .001$).

Faunal abundance

A total of 27 x 30-minute video ‘samples’ were collected from within the seagrass meadows. A total of 85 faunal individuals (based on MaxN) from 12 different taxa were

recorded, of which 1 was a Cephalopod. Some individuals could only be identified to family level (e.g. Gobiidae and Lutjanidae). Total relative faunal abundance (MaxN) per sample ranged from 17 individuals at Bo Cho to 0 individuals (at all sites). The average relative fish abundance (MaxN) across all sites and samples was 1.7 ± 3.7 . In Taw Wet North this was 0.3 ± 0.6 , in Lampi East 3.0 ± 4.6 , in Bo Cho 2.9 ± 5.6 and Nyaung Wee was 0.8 ± 1.4 (see Fig. 3).

Species diversity

The mean number of species was highest at Lampi East, with 1.3 ± 2.1 species per sample. At Bo Cho, this was 1.0 ± 1.5 , and at Nyaung Wee this was 0.3 ± 0.5 . The mean number of species was lowest at Taw Wet North, with 0.2 ± 0.4 . Average sample diversity (Shannon Wiener H') was highest at Lampi East (0.3 ± 0.5) and Bo Cho (0.2 ± 0.4). There was no sample diversity at Nyaung Wee or Taw Wet North (0.0 ± 0.0) (Fig. 3).

The most abundant fish species across all sites were the northern whiting (*Sillago sihama*) (7.0 ± 4.5), the common silver-biddy (*Gerres oyena*) (3.1 ± 1.6) and the pearly-spotted wrasse (*Halichoeres bicolor*) (2.0 ± 1.2). While seven individuals of the seagrass wrasse (*Novaculoides macrolepidotus*) were recorded, these were observed in only one sample. *G. oyena* was most frequent across all sites, occurring in 14 % of samples, followed by *H. bicolor* (10 %) and individuals from the Gobiidae family (10 %) then *S. sihama* (8 %). In total, only two taxa were recorded at Taw Wet North and Nyaung Wee. Nine taxa were recorded at Lampi East and seven at Bo Cho. The most frequently observed species in Taw Wet North was *S. sihama*, which was present in 13 % of the samples. In Lampi East the most frequently observed species were *G. oyena* (25 %), fish from the Gobiidae family (25 %) and *H. bicolor* (17 %). The most frequently observed species at Bo Cho sand were *H. bicolor* (17 %), thumbprint emperor (*Lethrinus harak*) (25%) and *S. sihama* (17%). *G. oyena* (25%) was the most frequently sampled fish in Nyaung Wee (Table. 3).

Species assemblages

The faunal species assemblages within the four seagrass meadows were not significantly different from each other (ANOSIM, $R = 0.04$, $P = 0.067$). Pairwise tests confirmed individual inter-site differences between Taw Wet North and Lampi East ($R = 0.09$, $P < 0.05$). No grouping existed for samples from the four sites, indicating some over-lapping of species assemblages (Fig. 4).

Discussion

Despite historical reports of rich and abundant seagrass meadows in Myanmar (Soe-Htun et al. (2001)) the present study provides convincing evidence that the seagrass fisheries within the Myeik Archipelago are in a potentially perilous state. Evidence of the limited seagrass fishery adds to a growing literature suggesting that Myanmar's marine habitats are in decline (Howard et al., 2014; Russell, 2015, 2016). Although the absence of seagrass associated fauna is of concern, our study suggests that the floral component of these seagrass meadows is in a healthy state with minimal visible impacts from eutrophication or sedimentation.

Across the sites surveyed, only 12 taxa of motile fauna were recorded. Relative to other regional and global studies this is extremely low (Esteban et al., 2017; Unsworth et al., 2014a). Across the Indo-Pacific, the number of seagrass-associated fish species is high. Nearly 700 species of fish are reported to have been observed in seagrass meadows across the Indo Pacific, the most common species' being *Lethrinus harak*, *Siganus canaliculatus* and *Gerres oyena* (Unsworth et al., 2014a). The apparent lack of these species' in the present study is cause alone for concern.

Multiple studies from the Indo-Pacific region suggest that many recognised and important reef dwellers, for example, Lethrinids and Siganids, utilise multiple habitat types, yet these families were also sparse in the seagrass meadows sampled in the present study. One *Siganus canaliculatus* individual was observed at Lampi East, one *Lethrinus variagatus* individual was observed at Taw Wet North and one *Lethrinus harak* individual was observed at Bo Cho. While some seagrass dependant (known to spend their whole life in seagrass) species such as *Gerres oyena* were present (Berkstrom et al., 2013; Dorenbosch et al., 2005; Unsworth et al., 2008), their low abundance appears uncharacteristic of the Indo-Pacific region. Multiple habitats use by marine fauna is primarily related to foraging migrations (as adults) or ontogenetic dietary shifts (as juveniles) (Nagelkerken, 2009). Reliance on multiple habitats underlines the importance of all habitat types and connectivity for maintaining fish assemblages. It could, therefore, be the case that other connected supporting habitats within the study region are in poor health. Dynamite fishing is ripe within the Myeik archipelago, and formal enforcement measures to reduce/prevent this activity have only come into force in recent years within the Lampi Marine National Park (MNP) (MOECA and Oikos, 2015). However, despite new 'written' protection measures, which is a progressive step, dynamite continues to be used. However, coral reef habitats within the archipelago are of average-good condition based on the scale used by Habibi et al. (2007); (Howard et al., 2014; Obura et al.,

237 2014). Mangrove communities, although minor regarding extension (notably within Lampi
238 MNP), are also in good condition with recognised high ecological value (Oikos and BANCA,
239 2011). This suggests that the fishery resource is simple highly overexploited.

240 Seagrass meadows are well known to fishers in Myanmar as an important fishing and
241 invertebrate gleaning (e.g. sea cucumbers) area (Schneider et al., 2014). Local people often
242 refer to seagrass as *Leik-Sar-Phat-Myet*, (directly translated as.....) recognising it as the food
243 of marine turtles (Soe-Htun et al., 2015; Soe-Htun et al., 2001). So although seagrasses and
244 associated habitats within the Myeik Archipelago are remote from large human populations,
245 they are facing the common problems associated with extensive overfishing (even within an
246 artisanal small-scale fishery) and perhaps overgrazing seen across the Indo-Pacific region in
247 previous decades (McManus, 1997). Barrier Net fishing, with nets that close off entire bays
248 (Plate 1) are common, and trawlers operate close to shore targeting shrimps and other fish
249 species (Soe-Htun et al., 2015). Anecdotal field observations confirm these fishing practices
250 (table. 4) and confirm the removal of top predators, including sharks. The lack of top predatory
251 fish is a distressing finding highlighted by the present study and previous studies in connected
252 habitats (Howard et al., 2014; Russell, 2016). While we appreciate all samples from the present
253 study were collected during the day, and lower abundances of predatory fish can be expected
254 due to diel differences in feeding activity (Unsworth et al., 2007), there were no fish from
255 predatory fish families such as Carangidae, Serranidae or Lutjanidae recorded. Methods were
256 the same as those conducted in other seagrass meadows from the region where much higher
257 abundances were recorded (Esteban et al., 2017; Unsworth et al., 2015). Even in extremely low
258 abundance, these predatory fish are much more receptive when using daytime baited cameras.
259 The lack of predatory species is symptomatic of a highly exploited fishery (see Plate 1.).

260 Herbivorous fish species are well recognised for their key functional role in supporting
261 the resilience of tropical marine systems by consuming excess algal or epiphytic growth.
262 Essentially herbivorous fish species prevent tropical marine systems from flipping into a state
263 of algal dominance (Maxwell et al., 2017). In the present study, seagrass was generally healthy
264 across all four sites, with low algal and epiphyte cover suggesting that there may be more
265 herbivores present than the study observed. However, other environmental conditions for
266 seagrass growth were favourable (good water clarity, limited physical disturbance) which may
267 negate the immediate need for herbivores in this context and herbivores have also been
268 confirmed lacking in adjacent habitats (Howard et al., 2014; Obura et al., 2014). The lack of
269 herbivores remains a concern as it places the long-term resilience of seagrass within this
270 archipelago in doubt (Burkholder et al., 2013).

The present study provides an abundance and diversity baseline for seagrass meadows within the Myeik Archipelago; it also offers some optimism for the future. Sites with the highest fish abundance and diversity (Lampi East and Bo Cho) are within Myanmar's only Marine National Park (MOECAP and Oikos, 2015). It is possible that the recent development of a five-year management plan and on the ground support to ranger patrols by the international NGO Istituto Oikos is having a positive impact on the marine environment.

In conclusion, this study provides evidence of the extreme over-exploitation of seagrass meadows, and associated habitats within the Myeik Archipelago. Overexploitation is likely a result of the several thousand fishing boats that operate within the region with limited regulation coupled with the historical and frequent use of dynamite fishing (Howard et al., 2014; Obura et al., 2014; Russell, 2015, 2016; Soe-Htun et al., 2015). While seagrass meadows currently appear healthy their future is questionable given that poor land-use management is resulting in an increase in land clearing for increased agriculture. The altered marine food web (lacking top predators) is likely damaging to the long-term resilience of Myanmar's seagrass ecosystems. Many of the fish species that would have been expected to be seen and are known to be important for food and coastal livelihoods throughout the region, particularly the Emperors, Rabbitfish and the Snappers (Unsworth and Cullen 2010), were absent. Although overexploited, the fish assemblages of coral areas will continue to provide some form of food supply to local people. However, the lack of fish within seagrass meadows here (thus a lack of juvenile recruitment), and the lack of predatory fish species means this food supply is in doubt for future. Current steps are, however, underway to include more seagrass sites within an MPA network planned for the archipelago with Taw Wet included in one of three large protected area sites currently being nominated for MPA status (Dearden, 2016). Overall, the evidence presented here suggests fisheries management is urgently and drastically required to bring sustainability to supporting seagrass stocks.

Acknowledgements

The present study was conducted with funding support from the Sustainable Places Research Institute and logistical support from Fauna & Flora International (Myanmar). The authors would like to thank Salai Mon Nyi Nyi Lin and Antt Maung for their logistical assistance in the field.

References

305 Beck, M.W., Heck Jr, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern,
306 B., Hays, C.G., Hoshino, K., Minello, T.J., 2001. The identification, conservation, and
307 management of estuarine and marine nurseries for fish and invertebrates. *BioScience* 51,
308 633-641.

309 Berkstrom, C., Lindborg, R., Thyresson, M., Gullstrom, M., 2013. Assessing connectivity in a
310 tropical embayment: Fish migrations and seascape ecology. *Biological Conservation* 166, 43-
311 53.

312 Burke, L., Reyntar, K., Spalding, M., Perry, A., 2011. *Reefs at Risk Revisited*. World Resources
313 Institute, Washington, DC, p. 114.

314 Burkholder, D.A., Heithaus, M.R., Fourqurean, J.W., Wirsing, A., Dill, L.M., 2013. Patterns of
315 top-down control in a seagrass ecosystem: could a roving apex predator induce a behaviour-
316 mediated trophic cascade? *Journal of Animal Ecology* 82, 1192-1202.

317 Cappel, M., Speare, P., De'ath, G., 2004. Comparison of baited remote underwater video
318 stations (BRUVS) and prawn (shrimp) trawls for assessments of fish biodiversity in inter-reefal
319 areas of the Great Barrier Reef Marine Park. *J. Exp. Mar. Biol. Ecol.* 302, 123-152.

320 Clarke, K.R., Warwick, R.M., 1994. *Changes in marine communities: an approach to statistical*
321 *analysis and interpretation*. Natural Environmental Research Council, Plymouth Marine
322 Laboratory, Plymouth, United Kingdom.

323 de la Torre-Castro, M., Di Carlo, G., Jiddawi, N.S., 2014. Seagrass importance for a small-scale
324 fishery in the tropics: The need for seascape management. *Mar. Poll. Bull.* 83, 398-407.

325 de la Torre-Castro, M., Ronnback, P., 2004. Links between humans and seagrasses - an
326 example from tropical East Africa. *Ocean & Coastal Management* 47, 361-387.

327 Dearden, P., 2016. *Blueprint for a Network of Marine Protected Areas in the Myeik*
328 *Archipelago, Myanmar*. Report No.39 of the Tanintharyi Conservation Programme, a joint
329 initiative of Fauna & Flora International (FFI), Myanmar Department of Fisheries and the
330 Myanmar Forest Department. FFI, Yangon.

331 Donner, S.D., Potere, D., 2007. The Inequity of the Global Threat to Coral Reefs. *BioScience*
332 57, 214-215.

333 Dorenbosch, M., Grol, M.G.G., Nagelkerken, I., van der Velde, G., 2005. Distribution of coral
334 reef fishes along a coral reef-seagrass gradient: edge effects and habitat segregation. *Mar.*
335 *Ecol. Prog. Ser.* 299, 277-288.

336 Duarte, C.M., Dennison, W.C., Orth, R.J.W., Carruthers, T.J.B., 2008. The charisma of coastal
337 ecosystems: Addressing the imbalance. *Estuaries And Coasts* 31, 233-238.

338 Esteban, N.E., Unsworth, R.K.F., Gourlay, J.B.Q., Hays, G.C., 2017. A new baseline for tropical
339 seagrass: Deep-water meadows in a pristine Indian Ocean wilderness. In Review.

340 Exton, D.A., 2010. Nearshore fisheries of the Wakatobi, in: Clifton, J., Unsworth, R.K.F. (Eds.),
341 *Marine conservation and research in the Coral Triangle: the Wakatobi Marine National Park*.
342 Nova Scientific, New York.

343 Folke, C., 2006. Resilience: The emergence of a perspective for social-ecological systems
344 analyses. *Global Environmental Change* 16, 253-267.

345 Giardino, C., Bresciani, M., Fava, F., Matta, E., Brando, V.E., Colombo, R., 2016. Mapping
346 Submerged Habitats and Mangroves of Lampi Island Marine National Park (Myanmar) from
347 in Situ and Satellite Observations. *Remote Sens.* 8, 13.

348 Gillanders, B.M., 2006. Seagrasses, fish, and fisheries, in: Larkum, A.W., Orth, R.J., Duarte,
349 C.M. (Eds.), *Seagrasses: Biology, Ecology and Conservation*. Springer, pp. 503-536.

350 Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J., Duke, N., 2011.
 351 Status and distribution of mangrove forests of the world using earth observation satellite
 352 data. *Global Ecology and Biogeography* 20, 154-159.
 353 Habibi, A., Setiasih, N., Sartin, J., 2007. A decade of reef check monitoring: Indonesian coral
 354 reefs, condition and trends. *The Indonesian Reef Check Network* 32.
 355 Haggitt, T., Freeman, D., Lily, C., 2014. Baited Remote Underwater Video Guidelines.
 356 Harborne, A., Mumby, P., Micheli, F., Perry, C., Dahlgren, C., Holmes, K., Brumbaugh, D., 2006.
 357 The functional value of Caribbean coral reef, seagrass and mangrove habitats to ecosystem
 358 processes. *ADV MAR BIOL* 50, 57-189.
 359 Harborne, A.R., Mumby, P.J., Kappel, C.V., Dahlgren, C.P., Micheli, F., Holmes, K.E.,
 360 Brumbaugh, D.R., 2008. Tropical coastal habitats as surrogates of fish community structure,
 361 grazing, and fisheries value. *Ecol. Applic.* 18, 1689-1701.
 362 Heck, K., Hays, G., Orth, R., 2003. Critical evaluation of the nursery role hypothesis for
 363 seagrass meadows. *Mar. Ecol. Prog. Ser.* 253, 123-136.
 364 Howard, R., Lunn, Z., Maung, A., Mon, S., Nyi Nyi, L., Thith, S., Aung, S., 2014. Assessment of
 365 the Myeik Archipelago Coral Reef Ecosystem, Reef Check Surveys, January 2013 to May 2014.
 366 Report No. 5 of the Tanintharyi Conservation Programme, a joint initiative of Fauna & Flora
 367 International (FFI) and the Myanmar Forest Department. FFI, Yangon.
 368 Ilangakoon, A.D., Tun, T., 2007. Rediscovering the Dugong (*Dugong dugon*) in myanmar and
 369 capacity building for research and conservation. *Raffles Bulletin of Zoology* 55, 195-199.
 370 Kelahe, B.P., Coleman, M.A., Broad, A., Rees, M.J., Jordan, A., Davis, A.R., 2014. Changes in
 371 fish assemblages following the establishment of a network of no-take marine reserves and
 372 partially-protected areas. *PloS one* 9, e85825.
 373 Malcolm, H.A., Schultz, A.L., Sachs, P., Johnstone, N., Jordan, A., 2015. Decadal changes in the
 374 abundance and length of snapper (*Chrysophrys auratus*) in subtropical marine sanctuaries.
 375 *PloS one* 10, e0127616.
 376 Maxwell, P.S., Eklöf, J.S., van Katwijk, M.M., O'Brien, K.R., de la Torre-Castro, M., Boström, C.,
 377 Bouma, T.J., Krause-Jensen, D., Unsworth, R.K.F., van Tussenbroek, B.I., van der Heide, T.,
 378 2017. The fundamental role of ecological feedback mechanisms for the adaptive
 379 management of seagrass ecosystems – a review. *Biological Reviews* 92, 1521-1538.
 380 McClanahan, T.R., Castilla, J.C., White, A.T., Defeo, O., 2009. Healing small-scale fisheries by
 381 facilitating complex socio-ecological systems. *Reviews in Fish Biology and Fisheries* 19, 33-47.
 382 McKenzie, L.J., Campbell, S.J., Roder, C.A., 2001. Seagrass-Watch: Manual for Mapping &
 383 Monitoring Seagrass Resources by Community (citizen) volunteers. Queensland Fisheries
 384 Service, Department of Primary Industries, Cairns, p. 100pp.
 385 McManus, J.W., 1997. Tropical marine fisheries and the future of coral reefs: a brief review
 386 with emphasis on Southeast Asia. *Coral Reefs* 16, S121-S127.
 387 MOECA, Oikos, 2015. Lampi Marine National Park: General Management Plan. 2014-2018.
 388 Ministry of Environmental Conservation and Forestry (MOECA), Yangon, Myanmar &
 389 Instituto Oikos, Milano, Italy.
 390 Nagelkerken, I., 2009. Ecological connectivity among Tropical Coastal Ecosystems Springer,
 391 Dordrecht.
 392 Nagelkerken, I., Blaber, S.J.M., Bouillon, S., Green, P., Haywood, M., Kirton, L.G., Meynecke,
 393 J.O., Pawlik, J., Penrose, H.M., Sasekumar, A., Somerfield, P.J., 2008. The habitat function of
 394 mangroves for terrestrial and marine fauna: A review. *Aquat. Bot.* 89, 155-185.
 395 Nagelkerken, I., Grol, M.G.G., Mumby, P.J., 2012. Effects of Marine Reserves versus Nursery
 396 Habitat Availability on Structure of Reef Fish Communities. *Plos One* 7, e36906.

397 Nordlund, L.M., Unsworth, R.K.F., Gullström, M., Cullen-Unsworth, L.C., 2017. Global
 398 significance of seagrass fishery activity. *Fish and Fisheries*, n/a-n/a.
 399 Novak, A.B., Hines, E., Kwan, D., Parr, L., Tun, M.T., Win, H., Short, F.T., 2009. Revised ranges
 400 of seagrass species in the Myeik Archipelago, Myanmar. *Aquat. Bot.* 91, 250-252.
 401 Obura, D.O., Benbow, S., Lunn, Z., 2014. Coral Diversity And Reef Resilience In The Northern
 402 Myeik Archipelago, Myanmar. Report No. 3 of the Tanintharyi Conservation Programme, a
 403 joint initiative of Fauna and Flora International (FFI) and the Myanmar Forest Department.
 404 FFI, Yangon.
 405 Oikos, BANCA, 2011. Myanmar Protected Areas: Context, Surrent Status and Challenges.
 406 Institutio Oikos, Milano, Italy & Biodiversity And Nature Conservation Association (BANCA),
 407 Yangon, Myanmar, Milano, Italy: Ancora Libri.
 408 Ooi, J.L.S., Kendrick, G.A., Van Niel, K.P., Affendi, Y.A., 2011. Knowledge gaps in tropical
 409 Southeast Asian seagrass systems. *Est. Coast. Shelf Sci.* 92, 118-131.
 410 Pauly, D., 1998. Tropical fishes: patterns and propensities. *J. Fish Biol.*, 1-17.
 411 Priede, I.G., Bagley, P.M., Smith, A., Creasey, S., Merrett, N.R., 1994. Scavenging deep
 412 demersal fishes of the porcupine seabight, northeast atlantic - observations by baited camera,
 413 trap and trawl. *Journal Of The Marine Biological Association Of The United Kingdom* 74, 481-
 414 498.
 415 Russell, B.C., 2015. Survey of coral reef fishes of the Myeik Archipelago, Myanmar. Report No.
 416 13 of the Tanintharyi Conservation Programme, a joint initiative of Fauna & Flora
 417 International (FFI) and the Myanmar Forest Department, FFI, Yangon, and the Bay of Bengal
 418 Large Marine Ecosystem project (BOBLME).
 419 Russell, B.C., 2016. Survey of coral reef fishes of the Myeik Archipelago, Myanmar. Report No.
 420 38 of the Tanintharyi Conservation Programme, a joint initiative of Fauna & Flora
 421 International (FFI) and the Myanmar Forest and Fisheries Departments. FFI, Yangon.
 422 Satumanatpan, S., 2008. National strategic and action plan for management seagrasses and
 423 dugong., in: Satumanatpan, S. (Ed.), Department of marine and coastal resources, Thailand.
 424 Satumanatpan, S., Thummikakpong, S., Kanongdate, K., 2011. Biodiversity of benthic fauna in
 425 the seagrass ecosystem of Kung Krabaen Bay, Chantaburi Province, Thailand. *Songklanakarin*
 426 *Journal of Science & Technology* 33.
 427 Schneider, H., Thiha, S., Pontillas, M.D., de Leon, E.M.C.P., 2014. Socio-economic baseline
 428 assessment: Thayawthatangyi and Langann Islands, Myeik Archipelago, Myanmar. Report No.
 429 10 of the Tanintharyi Conservation Programme, a joint initiative of Fauna & Flora
 430 International (FFI) and the Myanmar Forest Department, FFI, Yangon, and the Bay of Bengal
 431 Large Marine Ecosystem project (BOBLME).
 432 Soe-Htun, U., Maung, A., Mon, S., Ha, S., Aung, S., Lwin, A., Lunn, Z., 2015. Seagrass
 433 Conservation and Monitoring in Myanmar: The biodiversity, distribution and coverage of
 434 seagrasses in the Tanintharyi and Rakhine. Report No. 26 of the Tanintharyi Conservation
 435 Programme, a joint initiative of Fauna & Flora International (FFI) and the Myanmar Fisheries
 436 Department, FFI, Yangon and the Bay of Bengal Large Marine Ecosystem Project (BOBLME). .
 437 Soe-Htun, U., San-Tha-Htun, U., Mu-Mu-Aye, D., Ni-ni-win, D., Lei-lei-win, D., Ohno, M., 2001.
 438 Notes on seagrasses along Myanmar coastal regions. *Bull. Mar. Sci. Fish., Kochi Univ.* 21, 13-
 439 22.
 440 Unsworth, R.K.F., Cullen, L.C., 2010. Recognising the necessity for Indo-Pacific seagrass
 441 conservation. *Conservation Letters* 3, 63-73.

Unsworth, R.K.F., De Leon, P.S., Garrard, S.L., Jompa, J., Smith, D.J., Bell, J.J., 2008. High connectivity of Indo-Pacific seagrass fish assemblages with mangrove and coral reef habitats. *Marine Ecology-Progress Series* 353, 213-224.

Unsworth, R.K.F., Hinder, S.L., Bodger, O.G., Cullen-Unsworth, L.C., 2014a. Food supply depends on seagrass meadows in the coral triangle. *Environmental Research Letters* 9, 094005.

Unsworth, R.K.F., Jones, B.J., West, A., 2015. Baseline assessment of fish assemblages of Palma Bay, Mozambique. Report for MacAlister Elliott and Partners Ltd (MEP) Worldwide Fishery Consultants.

Unsworth, R.K.F., Peters, J.R., McCloskey, R.M., Hinder, S.L., 2014b. Optimising stereo baited underwater video for sampling fish and invertebrates in temperate coastal habitats. *Estuarine Coastal And Shelf Science* 150, 281–287.

Unsworth, R.K.F., Wylie, E., Smith, D.J., Bell, J.J., 2007. Diel trophic structuring of seagrass bed fish assemblages in the Wakatobi Marine National Park, Indonesia. *Estuarine Coastal And Shelf Science* 72, 81-88.

Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Short, F.T., Williams, S.L., 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* 106, 12377-12381.

Wraith, J., Lynch, T., Minchinton, T.E., Broad, A., Davis, A.R., 2013. Bait type affects fish assemblages and feeding guilds observed at baited remote underwater video stations. *Mar. Ecol. Prog. Ser.* 477, 189-199.

Wraith, J.A., 2007. Assessing reef fish assemblages in a temperate marine park using baited remote underwater video, School of Biological Sciences. University of Wollongong.

Table 1. Description of the four seagrass sites within the Myeik Archipelago, their designation, proximity to population and adjacent habitats.

Site	Designation	Population	Proximity of nearest population	Adjacent habitat	
				<i>mangrove</i>	<i>coral reef</i>
Taw Wet North (Taw Wet I.)	None	Uninhabited	~ 8.50 km	✓	*
Lampi East (Lampi I.)	Marine National Park	Inhabited	~ 3.25 km	✗	*
Bo Cho I.	Marine National Park	Inhabited	~ 0.05 km	✗	*
Nyaung Wee I.	None	Inhabited	~ 1 km	✓	✓

* While coral reef was present on-site observations confirmed that this was degraded and characteristic of previous bomb fishing.

Table 2. Seagrass cover and species composition at four sites within the Myeik Archipelago where *Cr* = *Cymodocea rotundata*, *CS* = *C. serrulata*, *Ho* = *Halophila ovalis*, *Hu* = *Halodule uninervis* and *Th* = *Thalassia hemrichii*.

Location	% Cover	<i>Cr</i>	<i>Cs</i>	<i>Ho</i>	<i>Hu</i>	<i>Th</i>
Taw Wet	32.1 ± 25.5	95.4 ± 11.6	3.3 ± 11.4	1.2 ± 2.7	0.00	0.00
Lampi East	42.6 ± 28.7	93.5 ± 14.3	0.3 ± 1.8	6.0 ± 13.7	0.9 ± 3.4	0.5 ± 2.0
Bo Cho	28.4 ± 21.1	85.1 ± 18.8	17.5 ± 15.1	0.0	0.0	0.0
Nyaung Wee	48.9 ± 27.5	93.8 ± 13.9	0.0	1.4 ± 6.5	4.8 ± 12.9	0.0

Table 3. Presence of individual species of fish recorded in samples using mono Baited Remote Underwater Video systems from four sites across the Myeik Archipelago, as a percentage of the total number of samples from each site.

Family	Species	Common Name	Taw Wet North	Lampi East	Bo Cho	Nyaung Wee
Gerreidae	<i>Gerres oyena</i>	Common silver bidy	-	25	8	25
Gobiidae		Goby	-	25	8	8
Labridae	<i>Halichoeres bicolor</i>	Pearly-spotted wrasse	-	17	25	-
Labridae	<i>Novaculoides macrolepidotus</i>	Seagrass wrasse	-	8	-	-
Lethrinidae	<i>Lethrinus harak</i>	Thumbprint emperor	-	-	25	-
Lethrinidae	<i>Lethrinus variegatus</i>	Slender emperor	13	-	-	-
Lutjanidae			-	8	-	-
Mugilidae	<i>Chelon spp.</i>	Mullet	-	8	-	-
Mullidae	<i>Parupeneus barberinus</i>	Dash-and-dot goatfish	-	8	8	-
Pomacentridae	<i>Pomacentrus spp.</i>	Damselfishes	-	8	-	-
Siganidae	<i>Siganus canaliculatus</i>	White-spotted spinefoot	-	8	-	-
Sillaginidae	<i>Shillago siamma</i>	Northern whiting	7	8	17	-
Tetraodontidae	<i>Arothron hispidus</i>	White-spotted puffer	-	-	8	-

490 Table 4. One observational landing survey from Nyaung Wee, from Thai crab fishermen
491 operating a trawler on soft sedimentary coastal areas within the Myeik Archipelago.

492

Catch	Weight (kg)	Species observed
Invertebrates	24	Mud crab
Fish	60	20+ Blue spotted stingray (<i>Neotrygon kuhlii</i>) 5 + White spotted puffer (<i>Arothron hispidus</i>) 2 small sharks, (<i>Carharhinus spp.</i>). 30+ Spotted scat (<i>Scatophagus argus</i>)

493

494

495




Figure 1. The location of the Myeik Archipelago on the southern coast of Myanmar. Inset: biodiversity and seagrass survey locations in the Myeik Archipelago during April and May 2016.




Figure 2. Mean (\pm SD) seagrass cover, canopy height, epiphyte cover and algae cover for four seagrass meadows across the Myeik Archipelago, Myanmar during April and May 2016.




Figure 3. Mean (\pm SD) abundance and number of species of motile fauna recorded within four seagrass meadows across the Myeik Archipelago, Myanmar during April and May 2016 using mono Baited Remote Underwater Video systems.




Figure 4. Two-dimensional non-metric MDS scaling configuration for comparisons between motile faunal assemblages recorded within seagrass meadows at four locations during April and May 2016 the Myeik Archipelago, Myanmar using mono Baited Remote Underwater Video systems.



517



518

519 Plate 1. Fishers at Taw Wet North utilising a barrier net that stretches across the entire bay
520 (638 m), to collect fish as the tide retreats. Picture provided by Ko Htwe.

521